

## NUCLEAR AND PARTICLE PHYSICS

### NET/JRF (JUNE-2011)

Q1. The radius of a  ${}^{64}_{29}\text{Cu}$  nucleus is measured to be  $4.8 \times 10^{-13}$  cm.

(A). The radius of a  ${}^{27}_{12}\text{Mg}$  nucleus can be estimated to be

- (a)  $2.86 \times 10^{-13}$  cm      (b)  $5.2 \times 10^{-13}$  cm      (c)  $3.6 \times 10^{-13}$  cm      (d)  $8.6 \times 10^{-13}$  cm

Ans: (c)

Solution: Since  $R = R_0(A)^{1/3} \Rightarrow \frac{R_{\text{Mg}}}{R_{\text{Cu}}} = \left(\frac{A_{\text{Mg}}}{A_{\text{Cu}}}\right)^{1/3} = \left(\frac{27}{64}\right)^{1/3}$


$$\Rightarrow \frac{R_{\text{Mg}}}{R_{\text{Cu}}} = \frac{3}{4} \Rightarrow R_{\text{Mg}} = \frac{3}{4} \times 4.8 \times 10^{-13} = 3.6 \times 10^{-13} \text{ cm.}$$

(B). The root-mean-square (r.m.s) energy of a nucleon in a nucleus of atomic number  $A$  in its ground state varies as:

- (a)  $A^{4/3}$       (b)  $A^{1/3}$       (c)  $A^{-1/3}$       (d)  $A^{-2/3}$

Ans: (c)

Q2. A beam of pions ( $\pi^+$ ) is incident on a proton target, giving rise to the process



(A). Assuming that the decay proceeds through strong interactions, the total isospin  $I$  and its third component  $I_3$  for the decay products, are

- (a)  $I = \frac{3}{2}, I_3 = \frac{3}{2}$       (b)  $I = \frac{5}{2}, I_3 = \frac{5}{2}$   
 (c)  $I = \frac{5}{2}, I_3 = \frac{3}{2}$       (d)  $I = \frac{1}{2}, I_3 = -\frac{1}{2}$

Ans: (c)

Solution:  $\pi^+ + p \rightarrow n + \pi^+ + \pi^+$ ;  $I: \frac{1}{2} + 1 + 1 = \frac{5}{2}$ ,  $I_3: -\frac{1}{2} + 1 + 1 = \frac{3}{2}$

(B). Using isospin symmetry, the cross-section for the above process can be related to that of the process

- (a)  $\pi^- n \rightarrow p \pi^- \pi^-$       (b)  $\pi^- \bar{p} \rightarrow \bar{n} \pi^- \pi^-$   
 (c)  $\pi^+ n \rightarrow p \pi^+ \pi^-$       (d)  $\pi^+ \bar{p} \rightarrow n \pi^+ \pi^-$

Ans: (c)



**NET/JRF (JUNE-2012)**

- Q5. The ground state of  ${}_{12}^{207}\text{Pb}$  nucleus has spin-parity  $J^p = \frac{1}{2}^-$ , while the first excited state has  $J^p = \frac{5}{2}^-$ . The electromagnetic radiation emitted when the nucleus makes a transition from the first excited state to ground state are
- (a) E2 and E3                      (b) M2 or E3                      (c) E2 or M3                      (d) M2 or M3

Ans: (c)

Solution: No parity change;  $\Delta J = 2, 3$

For  $E_l$  type,  $\Delta\pi = (-1)^l$ , (for no parity change  $l = 2$ )

For  $M_l$  type,  $\Delta\pi = (-1)^{l+1}$ , (for no parity change  $l = 3$ )

$\Delta J = 2$ , No parity change  $\rightarrow E2$ ;  $\Delta J = 3$ , No parity change  $\rightarrow M3$

- Q6. The dominant interactions underlying the following processes
- A.  $K^- + p \rightarrow \Sigma^- + \pi^+$ , B.  $\mu^- + \mu^+ \rightarrow K^- + K^+$ , C.  $\Sigma^+ \rightarrow p + \pi^0$  are
- (a) A: strong, B: electromagnetic and; C: weak  
 (b) A: strong, B: weak and; C: weak  
 (c) A: weak, B: electromagnetic and; C: strong  
 (d) A: weak, B: electromagnetic and; C: weak

Ans: (a)

(A)  $K^- + p \rightarrow \Sigma^- + \pi^+$  (Strong interaction)

$$I_3 : -\frac{1}{2} + \frac{1}{2} \rightarrow -1 + 1 \text{ (Conserved)}$$

(B)  $\mu^- + \mu^+ \rightarrow K^- + K^+$  (Electromagnetic interaction)

(C)  $\Sigma^+ \rightarrow p + \pi^0$  (Weak interaction)

$$I_3 : 1 \rightarrow \frac{1}{2} + 0 \text{ (Not conserved)}$$

**NET/JRF (JUNE-2013)**

Q7. The binding energy of a light nucleus  $(Z, A)$  in MeV is given by the approximate formula

$$B(A, Z) \approx 16A - 20A^{2/3} - \frac{3}{4}Z^2A^{-1/3} + 30\frac{(N-Z)^2}{A}$$

where  $N = A - Z$  is the neutron number. The value of  $Z$  of the most stable isobar for a given  $A$  is

- (a)  $\frac{A}{2}\left(1 - \frac{A^{2/3}}{160}\right)^{-1}$       (b)  $\frac{A}{2}$       (c)  $\frac{A}{2}\left(1 - \frac{A^{2/3}}{120}\right)^{-1}$       (d)  $\frac{A}{2}\left(1 + \frac{A^{4/3}}{64}\right)^{-1}$

Ans: (a)

Solution:  $\left.\frac{\partial B}{\partial Z}\right|_{Z=Z'} = 0 \Rightarrow Z' = \frac{A}{2}\left(1 - \frac{A^{2/3}}{160}\right)^{-1}$

Q8. A spin-1/2 particle  $A$  undergoes the decay  $A \rightarrow B + C + D$ , where it is known that  $B$  and  $C$  are also spin-1/2 particles. The complete set of allowed values of the spin of the particle  $D$  is

- (a)  $\frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}, 3, \dots$       (b) 0, 1      (c)  $\frac{1}{2}$  only      (d)  $\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \dots$

Ans: (c)

Solution: Spin of the left side and combined spin of the products must be same to conserve the spin angular momentum conservation law.

Q9. Muons are produced through the annihilation of particle  $a$  and its anti-particle, namely the process  $a + \bar{a} \rightarrow \mu^+ + \mu^-$ . A muon has a rest mass of  $105 \text{ MeV}/c^2$  and its proper life time is  $2\mu\text{s}$ . If the center of mass energy of the collision is  $2.1 \text{ GeV}$  in the laboratory frame that coincides with the center-of-mass frame, then the fraction of muons that will decay before they reach a detector placed  $6 \text{ km}$  away from the interaction point is

- (a)  $e^{-1}$       (b)  $1 - e^{-1}$       (c)  $1 - e^{-2}$       (d)  $e^{-10}$

Ans: (b)

Solution:  $N = N_0 e^{-\lambda t} \Rightarrow \frac{N}{N_0} = e^{-\lambda t} = e^{-\frac{t}{\tau}}$ , where  $\tau = 2 \times 10^{-6} \text{ s}$ ,  $\gamma = \frac{2.1}{105} \times 10^3 = 20$  and

$$t = \frac{6 \times 10^3}{3 \times 10^8} = 2 \times 10^{-5} \text{ sec. Thus } \frac{t}{\gamma\tau} = \frac{1}{2} \Rightarrow \frac{N}{N_0} = e^{-\frac{1}{2}} \approx 1 - e^{-1}.$$

**NET/JRF -(DEC-2013)**

Q10. The intrinsic electric dipole moment of a nucleus  ${}^A_Z X$

- (a) increases with  $Z$ , but independent of  $A$
- (b) decreases with  $Z$ , but independent of  $A$
- (c) is always zero
- (d) increases with  $Z$  and  $A$

Ans: (d)

Q11. According to the shell model, the total angular momentum (in units of  $\hbar$ ) and the parity of the ground state of the  ${}^7_3 Li$  nucleus is

- (a)  $\frac{3}{2}$  with negative parity
- (b)  $\frac{3}{2}$  with positive parity
- (c)  $\frac{1}{2}$  with positive parity
- (d)  $\frac{7}{2}$  with negative parity

Ans: (a)

Solution:  $Z = 3, N = 4$

For odd  $Z = 3; (s^2)_{1/2} (p^1)_{3/2} \Rightarrow j = 3/2, l = 1$  and parity  $= (-1)^l = -1$ .

**NET/JRF (JUNE-2014)**

Q12. The recently-discovered Higgs boson at the LHC experiment has a decay mode into a photon and a  $Z$  boson. If the rest masses of the Higgs and  $Z$  boson are  $125 \text{ GeV}/c^2$  and  $90 \text{ GeV}/c^2$  respectively, and the decaying Higgs particle is at rest, the energy of the photon will approximately be

- (a)  $35\sqrt{3} \text{ GeV}$
- (b)  $35 \text{ GeV}$
- (c)  $30 \text{ GeV}$
- (d)  $15 \text{ GeV}$

Ans: (c)

Solution: Assume H is symbol of Higgs boson,  $H \rightarrow Z + \gamma$

$$E_\gamma = \frac{E_H^2 - E_Z^2}{2E_H} = \frac{(125)^2 - (90)^2}{2 \times 125} = 30 \text{ GeV}$$

Q13. In a classical model, a scalar (spin-0) meson consists of a quark and an antiquark bound by a potential  $V(r) = ar + \frac{b}{r}$ , where  $a = 200 \text{ MeV fm}^{-1}$  and  $b = 100 \text{ MeV fm}$ . If the masses of the quark and antiquark are negligible, the mass of the meson can be estimated as approximately

- (a)  $141 \text{ MeV}/c^2$       (b)  $283 \text{ MeV}/c^2$       (c)  $353 \text{ MeV}/c^2$       (d)  $425 \text{ MeV}/c^2$

Ans: (b)

Solution: At equilibrium separation the potential is minimum, thus the equilibrium separation can be determined as

$$\left. \frac{dV(r)}{dr} \right|_{r=r_0} = a - \frac{b}{r_0^2} = 0 \Rightarrow r_0 = \sqrt{\frac{b}{a}} = \sqrt{\frac{100 \text{ MeV fm}}{200 \text{ MeV fm}^{-1}}} = \frac{1}{\sqrt{2}} \text{ fm}$$

The equilibrium separation between particles is also estimated by uncertainty principle

$$r_0 = c\Delta t \Rightarrow r_0 = c \frac{\hbar}{\Delta E} \quad (\text{where, } \Delta E \Delta t \approx \hbar)$$

Where,  $c$  is the velocity of the virtual meson

$$r_0 = c \frac{\hbar}{\Delta E} = \frac{200 \text{ MeV} \cdot \text{fm}}{\Delta E (\text{MeV})}$$

Using above two relation  $\frac{200 \text{ MeV} \cdot \text{fm}}{\Delta E (\text{MeV})} = \frac{1}{\sqrt{2}} \text{ fm}$

$$\Delta E = 200\sqrt{2} = 283 \text{ MeV} \Rightarrow \Delta E = \Delta m \times c^2$$

the mass of the meson  $\Delta m = \frac{\Delta E}{c^2} = 283 \text{ MeV} / c^2$

### NET/JRF (DEC-2014)

Q14. Consider the four processes

- (i)  $p^+ \rightarrow n + e^+ + \nu_e$       (ii)  $\Lambda^0 \rightarrow p^+ + e^+ + \nu_e$   
 (iii)  $\pi^+ \rightarrow e^+ + \nu_e$       (iv)  $\pi^0 \rightarrow \gamma + \gamma$

which of the above is/are forbidden for free particles?

- (a) only (ii)      (b) (ii) and (iv)      (c) (i) and (iv)      (d) (i) and (ii)

Ans: (d)

Solution: (i)  $p^+ \rightarrow n + e^+ + \nu_e$  [Not allowed]

It violate energy conservation. The mass of proton is less than mass of neutron. Free proton is stable and can not decay to neutron. Proton can decay to neutron only inside the nucleus, where energy violation is taken care by Heisenberg uncertainty principle.

(ii)  $\Lambda^0 \rightarrow p^+ + e^+ + \nu_e$  [Not allowed]. In this decay charge is not conserved

(iii)  $\pi^+ \rightarrow e^+ + \nu_e$  [allowed through Weak interaction]

(iv)  $\pi^0 \rightarrow \gamma + \gamma$  [allowed through Electromagnetic interaction]

Q15. In deep inelastic scattering electrons are scattered off protons to determine if a proton has any internal structure. The energy of the electron for this must be at least

- (a)  $1.25 \times 10^9 eV$       (b)  $1.25 \times 10^{12} eV$       (c)  $1.25 \times 10^6 eV$       (d)  $1.25 \times 10^8 eV$

Ans: (b)

Solution: The internal structure of proton can only be determined if the wavelength of the incoming electron is nearly equal to the size of the proton

i.e.  $\lambda = R = 1.2A^{1/3} (fm) = 1.2 fm = 1.2 \times 10^{-15} m$

According to de-Broglie relation,  $\lambda = \frac{h}{p} = \frac{h}{\sqrt{2mE}}$

This can be also written as  $\lambda \left( \overset{\circ}{\text{A}} \right) = \sqrt{\frac{150}{E(eV)}}$

$$\therefore E(eV) = \frac{150}{\left[ \lambda \left( \overset{\circ}{\text{A}} \right) \right]^2} = \frac{150}{(1.2 \times 10^{-5})^2} = 1.04 \times 10^{12} \Rightarrow E = 1.04 \times 10^{12} eV$$

The bet suitable answer is option (b).

Q16. If the binding energy  $B$  of a nucleus (mass number  $A$  and charge  $Z$ ) is given by

$$B = a_v A - a_s A^{2/3} - a_{sym} \frac{(2Z - A)^2}{A} + \frac{a_c Z^2}{A^{1/3}}$$

where  $a_v = 16 MeV$ ,  $a_s = 16 MeV$ ,  $a_{sym} = 24 MeV$  and  $a_c = 0.75 MeV$ , then for the most stable isobar for a nucleus with  $A = 216$  is

- (a) 68                      (b) 72                      (c) 84                      (d) 92

Ans: (c)

Solution: For the most stable isobar for a nucleus  $\frac{dB}{dZ} = 0 \Rightarrow -a_{sym} \frac{2(2Z - A) \times 2}{A} + \frac{2a_c Z}{A^{1/3}} = 0$

$$\Rightarrow 24 \frac{2(2Z - 216) \times 2}{216} + 0.75 \frac{2Z}{(216)^{1/3}} = 0 \Rightarrow \frac{4(2Z - 216)}{9} + \frac{3 \cdot 2Z}{4 \cdot 6} = 0$$

$$\Rightarrow \frac{4(2Z - 216)}{9} + \frac{Z}{4} = 0 \Rightarrow 16(2Z - 216) + 9Z = 0 \Rightarrow 41Z = 216 \times 16 \Rightarrow Z = 82.3$$

### NET/JRF (JUNE-2015)

Q17. The reaction  ${}^2_1D + {}^2_1D \rightarrow {}^4_2He + \pi^0$  cannot proceed via strong interactions because it violates the conservation of

- (a) angular momentum (b) electric charge  
(c) baryon number (d) isospin

Ans. (d)

Solution:  ${}_1D^2 + {}_1D^2 \rightarrow {}_2He^4 + \pi^0$  (Not conserved)

$$I: 0 \quad 0 \rightarrow 0 \quad 1$$

This isospin is not conserved in above reaction.

Q18. Let us approximate the nuclear potential in the shell model by a three dimensional isotropic harmonic oscillator. Since the lowest two energy levels have angular momenta  $l=0$  and  $l=1$  respectively, which of the following two nuclei have magic numbers of protons and neutrons?

- (a)  ${}^4_2He$  and  ${}^{16}_8O$  (b)  ${}^2_1D$  and  ${}^8_4Be$  (c)  ${}^4_2He$  and  ${}^8_4Be$  (d)  ${}^4_2He$  and  ${}^{12}_6C$

Ans. (a)

Solution:  ${}_2He^4$  has  $Z=2, N=2$

and  ${}_8O^{16}$  has  $Z=8, N=8$  magic numbers (2, 8, 20, 28, 50, 82, 126)

Q19. The charm quark  $S$  assigned a charm quantum number  $C=1$ . How should the Gellmann-Nishijima formula for electric charge be modified for four flavors of quarks?

- (a)  $I_3 + \frac{1}{2}(B - S - C)$  (b)  $I_3 + \frac{1}{2}(B - S + C)$   
(c)  $I_3 + \frac{1}{2}(B + S - C)$  (d)  $I_3 + \frac{1}{2}(B + S + C)$







- Q23. In the large hadron collider (*LHC*), two equal energy proton beams traverse in opposite directions along a circular path of length  $27\text{ km}$ . If the total centre of mass energy of a proton-proton pair is  $14\text{ TeV}$ , which of the following is the best approximation for the proper time taken by a proton to traverse the entire path?
- (a)  $12\text{ ns}$                       (b)  $1.2\ \mu\text{s}$                       (c)  $1.2\text{ ns}$                       (d)  $0.12\ \mu\text{s}$

Ans: (a)

Solution: The proton travel at nearly speed of light in *LHC*, therefore

$$t \approx \frac{d}{c} = \frac{27 \times 10^3}{3 \times 10^8} \approx 9 \times 10^{-5}\text{ sec}$$

Since, proton is relativistic,  $t_0 = t \sqrt{1 - \frac{v^2}{c^2}} = \frac{t}{\gamma}$

$$\because E = \gamma m_0 c^2 \Rightarrow \frac{1}{\gamma} = \frac{m_0 c^2}{E} = \frac{938\text{ MeV}}{7\text{ TeV}} = \frac{938 \times 10^6\text{ eV}}{7 \times 10^{12}\text{ eV}} = 1.34 \times 10^{-4}$$

Thus,  $t_0 = \frac{t}{\gamma} = 9 \times 10^{-5} \times 1.34 \times 10^{-4} = 1.2 \times 10^{-8}\text{ sec} = 12\text{ ns}$

- Q24. Let  $E_s$  denotes the contribution of the surface energy per nucleon in the liquid drop model. The ratio  $E_s({}_{13}^{27}\text{Al}) : E_s({}_{30}^{64}\text{Zn})$  is
- (a) 2:3                      (b) 4:3                      (c) 5:3                      (d) 3:2

Ans: (b)

Solution:  $E_s = \frac{B}{A} = \frac{A^{\frac{2}{3}}}{A} \propto A^{-\frac{1}{3}} \Rightarrow \frac{E_s(\text{Al})}{E_s(\text{Zn})} = \frac{(27)^{\frac{1}{3}}}{(64)^{\frac{1}{3}}} = \frac{(64)^{\frac{1}{3}}}{(27)^{\frac{1}{3}}} = \frac{4}{3}$

- Q25. According to the shell model, the nuclear magnetic moment of the  ${}_{13}^{27}\text{Al}$  nucleus is (Given that for a proton  $g_l = 1$ ,  $g_s = 5.586$ , and for a neutron  $g_l = 0$ ,  $g_s = -3.826$ )
- (a)  $-1.913\ \mu_N$                       (b)  $14.414\ \mu_N$                       (c)  $4.793\ \mu_N$                       (d) 0

Ans: (c)

Solution:  ${}_{13}\text{Al}^{27} : Z = 13, N = 14$  for  $Z = 13, S_{1/2}^2, P_{3/2}^4, P_{1/2}^2, d_{5/2}^5 \Rightarrow j = \frac{5}{2}, l = 2$

Magnetic moment,  $\mu = \frac{1}{2} [2j - 1 + g_s] \mu_N = \frac{1}{2} \left[ 2 \times \frac{5}{2} - 1 + 5.586 \right] \mu_N \Rightarrow \mu = 4.793\ \mu_N$

**NET/JRF (DEC-2016)**

Q26. What should be the minimum energy of a photon for it to split an  $\alpha$ -particle at rest into a tritium and a proton?

(The masses of  ${}^4_2\text{He}$ ,  ${}^3_1\text{H}$  and  ${}^1_1\text{H}$  are  $4.0026\text{amu}$ ,  $3.0161\text{amu}$  and  $1.0073\text{amu}$  respectively, and  $1\text{amu} \approx 938\text{MeV}$ )

- (a)  $32.2\text{MeV}$                       (b)  $3\text{MeV}$                       (c)  $19.3\text{MeV}$                       (d)  $931.5\text{MeV}$

Ans. : (c)

Solution: From conservation of energy

$$E_\alpha + m_\alpha c^2 = m_{{}^3_1\text{H}} c^2 + m_{{}^1_1\text{H}} c^2$$

$$\text{or } E_\alpha = [m_{{}^3_1\text{H}} + m_{{}^1_1\text{H}} - m_\alpha] \times 938\text{MeV} = 19.5\text{MeV}$$

Q27. Which of the following reaction(s) is/are allowed by the conservation laws?

(i)  $\pi^+ + n \rightarrow \Lambda^0 + K^+$

(ii)  $\pi^- + p \rightarrow \Lambda^0 + K^0$

- (a) both (i) and (ii)                      (b) only (i)  
 (c) only (ii)                                      (d) neither (i) nor (ii)

Ans. : (a)

Solution: (i)  $\pi^+ + n \rightarrow \Lambda^0 + K^+$

$q: 1+0 \rightarrow 0+1$

$B: 0+1 \rightarrow 1+0$

$S: 0+0 \rightarrow -1+1$

Reaction is allowed

(ii)  $\pi^- + p \rightarrow \Lambda^0 + K^0$

$q: -1+1 \rightarrow 0+0$

$B: 0+1 \rightarrow 1+0$

$S: 0+0 \rightarrow -1+1$

Reaction is allowed

Q28. A particle, which is a composite state of three quarks  $u, d$  and  $s$ , has electric charge, spin and strangeness respectively, equal to

- (a)  $1, \frac{1}{2}, -1$       (b)  $0, 0, -1$       (c)  $0, \frac{1}{2}, -1$       (d)  $-1, -\frac{1}{2}, +1$

Ans. : (c)

Solution: charge, spin and strangeness of Quarks  $u, d$  &  $s$  are given as

	U	D	S	Total
Charge	$\frac{+2}{3}$	$\frac{-1}{3}$	$\frac{-1}{3}$	0
Spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$ or $\frac{3}{2}$
Strangeness	0	0	-1	-1

If a particle  $x$  is a composite of  $u, d$  &  $s$ , then net charge, spin and strangeness on  $x$  is  
net charge = 0

net spin =  $\frac{1}{2}$  or  $\frac{3}{2}$  and net strangeness = -1

### NET/JRF (JUNE-2017)

Q29. If in a spontaneous  $\alpha$  - decay of  ${}_{92}^{232}\text{U}$  at rest, the total energy released in the reaction is  $Q$ , then the energy carried by the  $\alpha$  - particle is

- (a)  $57Q/58$       (b)  $Q/57$       (c)  $Q/58$       (d)  $23Q/58$

Ans. : (a)

Solution: Energy carried by the  $\alpha$  - particle is

$$KE_{\alpha} = \left( \frac{A-4}{A} \right) Q = \frac{228}{232} Q = \frac{57}{58} Q$$

Q30. The range of the nuclear force between two nucleons due to the exchange of pions is  $1.40 \text{ fm}$ . If the mass of pion is  $140 \text{ MeV}/c^2$  and the mass of the rho-meson is  $770 \text{ MeV}/c^2$ , then the range of the force due to exchange of rho-mesons is

- (a)  $1.40 \text{ fm}$       (b)  $7.70 \text{ fm}$       (c)  $0.25 \text{ fm}$       (d)  $0.18 \text{ fm}$

Ans. : (c)

Solution: Range for nuclear force between nucleon will be  $R = c\Delta t = \frac{\hbar c}{mc^2}$  and  $\hbar c = 199 \text{ MeVfm}$

$$\Rightarrow R = \frac{199 \text{ MeVfm}}{770 \frac{\text{MeV}}{c^2} \times c^2} \approx 0.25 \text{ fm}$$

Q31. A baryon  $X$  decays by strong interaction as  $X \rightarrow \Sigma^+ + \pi^- + \pi^0$ , where  $\Sigma^+$  is a member of the isotriplet  $(\Sigma^+, \Sigma^0, \Sigma^-)$ . The third component  $I_3$  of the isospin of  $X$  is

- (a) 0                      (b) 1/2                      (c) 1                      (d) 3/2

Ans. : (a)

Solution:  $X = \Sigma^+ + \pi^- + \pi^0$

$$I_3 : \underbrace{1 \quad -1 \quad 0}$$

$\Rightarrow I_3$  for  $X$  is 0.

