

MATHEMATICAL PHYSICS SOLUTIONS**GATE-2010**

- Q1. Consider an anti-symmetric tensor P_{ij} with indices i and j running from 1 to 5. The number of independent components of the tensor is
- (a) 3 (b) 10 (c) 9 (d) 6

Ans: (b)

Solution: The number of independent components of the tensor

$$= \frac{1}{2}(N^2 - N) = \frac{1}{2}(25 - 5) = 10 \quad \therefore N = 5$$

- Q2. The value of the integral $\oint_C \frac{e^z \sin(z)}{z^2} dz$, where the contour C is the unit circle: $|z - 2| = 1$, is
- (a) $2\pi i$ (b) $4\pi i$ (c) πi (d) 0

Ans: (d)

Solution: Pole is at $z = 0$, Circle $|z - 2| = 1 \Rightarrow \oint_C \frac{e^z \sin z}{z^2} dz = 2\pi i \times 0 = 0$.

- Q3. The eigenvalues of the matrix $\begin{pmatrix} 2 & 3 & 0 \\ 3 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ are
- (a) 5, 2, -2 (b) -5, -1, -1 (c) 5, 1, -1 (d) -5, 1, 1

Ans: (c)

Solution: The characteristic equation of the matrix A , $|A - \lambda I| = 0$

$$\Rightarrow |A - \lambda I| = \begin{vmatrix} 2 - \lambda & 3 & 0 \\ 3 & 2 - \lambda & 0 \\ 0 & 0 & 1 - \lambda \end{vmatrix} = 0 \Rightarrow \lambda^3 - 5\lambda^2 - \lambda + 5 = 0 \Rightarrow \lambda = 5, 1, -1$$

- Q4. If $f(x) = \begin{cases} 0 & \text{for } x < 3, \\ x - 3 & \text{for } x \geq 3 \end{cases}$ then the Laplace transform of $f(x)$ is
- (a) $s^{-2}e^{sx}$ (b) s^2e^{sx} (c) s^{-2} (d) $s^{-2}e^{-sx}$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Ans: (d)

$$\text{Solution: } L\{f(x)\} = \int_0^{\infty} e^{-sx} f(x) dx = \int_0^3 e^{-sx} f(x) dx + \int_3^{\infty} e^{-sx} f(x) dx = \int_3^{\infty} (x-3)e^{-sx} dx$$

$$L\{f(x)\} = (x-3) \frac{e^{-sx}}{-s} \Big|_3^{\infty} - \int_3^{\infty} 1 \cdot \left(\frac{e^{-sx}}{-s} \right) dx = 0 - \frac{1}{s} \int_3^{\infty} e^{-sx} dx = \frac{1}{s} \left[\frac{e^{-sx}}{-s} \right]_3^{\infty} = s^{-2} e^{-sx}$$

Q5. The solution of the differential equation for $y(t)$: $\frac{d^2 y}{dt^2} - y = 2 \cosh(t)$, subject to the initial conditions $y(0)=0$ and $\left. \frac{dy}{dt} \right|_{t=0} = 0$, is

(a) $\frac{1}{2} \cosh(t) + t \sinh(t)$

(b) $-\sinh(t) + t \cosh(t)$

(c) $t \cosh(t)$

(d) $t \sinh(t)$

Ans: (d)

Solution: For C.F. $(D^2 - 1)y = 0 \Rightarrow m = \pm 1 \Rightarrow C.F. = C_1 e^t + C_2 e^{-t}$

$$P.I. = \frac{1}{D^2 - 1} (2 \cosh t) = \frac{1}{D^2 - 1} 2 \left(\frac{e^t + e^{-t}}{2} \right) = \frac{1}{D^2 - 1} (e^t) + \frac{1}{D^2 - 1} (e^{-t}) = \frac{t}{2} e^t + \frac{t}{2} (-e^{-t})$$

$$\Rightarrow y = C_1 e^t + C_2 e^{-t} + \frac{t}{2} e^t - \frac{t}{2} e^{-t} \Rightarrow y(0) = 0 \Rightarrow C_1 + C_2 = 0$$

$$\frac{dy}{dt} = C_1 e^t - C_2 e^{-t} + \frac{t}{2} e^t + \frac{1}{2} e^t + \frac{t}{2} e^{-t} - \frac{1}{2} e^{-t}$$

$$\left. \frac{dy}{dt} \right|_{t=0} = 0 \Rightarrow C_1 - C_2 + 0 + \frac{1}{2} + 0 - \frac{1}{2} = 0 \Rightarrow C_1 - C_2 = 0$$

Since $C_1 + C_2 = 0$ and $C_1 - C_2 = 0 \Rightarrow C_1 = 0, C_2 = 0$.

$$\text{Thus } \Rightarrow y = \frac{t}{2} e^t - \frac{t}{2} e^{-t} \Rightarrow y = t \sinh t$$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

GATE-2011

Q6. Two matrices A and B are said to be similar if $B = P^{-1}AP$ for some invertible matrix P . Which of the following statements is NOT TRUE?

- (a) $\text{Det}A = \text{Det}B$ (b) Trace of $A = \text{Trace of } B$
 (c) A and B have the same eigenvectors (d) A and B have the same eigenvalues

Ans: (c)

Solution: If A and P be square matrices of the same type and if P be invertible then matrices A and $B = P^{-1}AP$ have the same characteristic roots

Then $B - \lambda I = P^{-1}AP - P^{-1}\lambda I P = P^{-1}(A - \lambda I)P$ where I is identity matrix.

$$|B - \lambda I| = |P^{-1}(A - \lambda I)P| = |P^{-1}||A - \lambda I||P| = |A - \lambda I||P^{-1}||P| = |A - \lambda I||PP^{-1}| = |A - \lambda I|$$

Thus the matrices A and $B (= P^{-1}AP)$ have the same characteristic equation and hence characteristic roots of eigen values. Since the sum of the eigen values of a matrix and product of eigen values of a matrix is equal to the determinant of matrix hence third alternative is incorrect.

Q7. If a force \vec{F} is derivable from a potential function $V(r)$, where r is the distance from the origin of the coordinate system, it follows that

- (a) $\vec{\nabla} \times \vec{F} = 0$ (b) $\vec{\nabla} \cdot \vec{F} = 0$ (c) $\vec{\nabla} V = 0$ (d) $\nabla^2 V = 0$

Ans: (a)

Solution: Since \vec{F} is derivative from potential $V(r)$ and $\vec{F} = -\vec{\nabla}V(r)$

$$\Rightarrow \vec{\nabla} \times \vec{F} = -\vec{\nabla} \times (\vec{\nabla}V) = 0.$$

Q8. A 3×3 matrix has elements such that its trace is 11 and its determinant is 36. The eigenvalues of the matrix are all known to be positive integers. The largest eigenvalues of the matrix is

- (a) 18 (b) 12 (c) 9 (d) 6

Ans: (c)

Solution: We know that for any matrix

- The product of eigenvalues is equal to determinant of that matrix.

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

2. $\lambda_1 + \lambda_2 + \lambda_3 + \dots =$ Trace of matrix

$\lambda_1 + \lambda_2 + \lambda_3 = 11$ and $\lambda_1\lambda_2\lambda_3 = 36$. Hence the largest eigen value of the matrix is 9.

Q9. The unit vector normal to the surface $x^2 + y^2 - z = 1$ at the point $P(1, 1, 1)$ is

- (a) $\frac{\hat{i} + \hat{j} - \hat{k}}{\sqrt{3}}$ (b) $\frac{2\hat{i} + \hat{j} - \hat{k}}{\sqrt{6}}$ (c) $\frac{\hat{i} + 2\hat{j} - \hat{k}}{\sqrt{6}}$ (d) $\frac{2\hat{i} + 2\hat{j} - \hat{k}}{3}$

Ans: (d)


Solution: The equation of the system is $f(x, y, z) \equiv (x^2 + y^2 - z - 1) = 0$

The gradient of the above function is $\vec{\nabla}f = \left(\frac{\partial}{\partial x}\hat{i} + \frac{\partial}{\partial y}\hat{j} + \frac{\partial}{\partial z}\hat{k} \right) (x^2 + y^2 - z - 1)$
 $= 2x\hat{i} + 2y\hat{j} - \hat{k}$

Hence unit normal vector at $(1, 1, 1)$ $\frac{\vec{\nabla}f}{|\vec{\nabla}f|} = \frac{2\hat{i} + 2\hat{j} - \hat{k}}{3}$.

Q10. Consider a cylinder of height h and radius a , closed at both ends, centered at the origin.

Let $\vec{r} = \hat{i}x + \hat{j}y + \hat{k}z$ be the position vector and \hat{n} a unit vector normal to the surface. The surface integral $\int_S \vec{r} \cdot \hat{n} ds$ over the closed surface of the cylinder is



- (a) $2\pi a^2 (a + h)$ (b) $3\pi a^2 h$ (c) $2\pi a^2 h$ (d) zero

Ans: (b)

Solution: $\oint_S \vec{r} \cdot \hat{n} ds = \int_V (\vec{\nabla} \cdot \vec{r}) d\tau = 3 \int_V d\tau = 3\pi a^2 h$

Q11. The solutions to the differential equation $\frac{dy}{dx} = -\frac{x}{y+1}$ are a family of

- (a) circles with different radii
 (b) circles with different centres

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
 Near IIT, Hauz Khas, New Delhi-16
 Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
 28-B/6, Jia Sarai, Near IIT
 Hauz Khas, New Delhi-16

- (c) straight lines with different slopes
 (d) straight lines with different intercepts on the y -axis

Ans: (a)

$$\text{Solution: } \frac{dy}{dx} = -\frac{x}{y+1} \Rightarrow xdx + ydy + dy = 0 \Rightarrow \frac{x^2}{2} + \frac{y^2}{2} + y = C_1 \Rightarrow x^2 + y^2 + 2y = 2C_1$$

$$\Rightarrow (x-0)^2 + (y+1)^2 = 2C_1 + 1 = C$$

which is family of circles with different radii.

- Q12. Which of the following statements is **TRUE** for the function $f(z)$?
- (a) $f(z)$ is analytic everywhere in the complex plane
 (b) $f(z)$ has a zero at $z = \pi$
 (c) $f(z)$ has a pole of order 2 at $z = \pi$
 (d) $f(z)$ has a simple pole at $z = \pi$

Ans: (c)

Solution: Consider a function $f(z) = \frac{z \sin z}{(z - \pi)^2}$ of a complex variable z .

$$f(z) = \frac{z \sin z}{(z - \pi)^2} \text{ has a pole of order 2 at } z = \pi$$

- Q13. Consider a counterclockwise circular contour $|z|=1$ about the origin. The integral $\oint f(z)dz$ over this contour is

- (a) $-i\pi$ (b) zero (c) $i\pi$ (d) $2i\pi$

Ans: (b)

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
 Near IIT, Hauz Khas, New Delhi-16
 Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
 28-B/6, Jia Sarai, Near IIT
 Hauz Khas, New Delhi-16

GATE-2012

Q14. Identify the correct statement for the following vectors $\vec{a} = 3\hat{i} + 2\hat{j}$ and $\vec{b} = \hat{i} + 2\hat{j}$

- (a) The vectors \vec{a} and \vec{b} are linearly independent
- (b) The vectors \vec{a} and \vec{b} are linearly dependent
- (c) The vectors \vec{a} and \vec{b} are orthogonal
- (d) The vectors \vec{a} and \vec{b} are normalized

Ans: (a)

Solution: If $\vec{a} = 3\hat{i} + 2\hat{j}$, $\vec{b} = \hat{i} + 2\hat{j}$ are linearly dependent $\vec{a} + m\vec{b} = 0$, for some values of m but $3 + m = 0$ and $2 + 2m = 0$ do not have any solution. So they are linearly independent.

$\vec{a} \cdot \vec{b} \neq 0$ (Not orthogonal); $\vec{a} \times \vec{b} \neq 0$ (Not normalized)

Q15. The number of independent components of the symmetric tensor A_{ij} with indices $i, j = 1, 2, 3$ is

- (a) 1
- (b) 3
- (c) 6
- (d) 9

Ans: (c) For symmetric tensor

$$A_{ij} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix}$$

$\therefore A_{12} = A_{21}, A_{23} = A_{32}, A_{13} = A_{31}$, hence there are six independent components.

Q16. The eigenvalues of the matrix $\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$ are

- (a) 0, 1, 1
- (b) $0, -\sqrt{2}, \sqrt{2}$
- (c) $\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0$
- (d) $\sqrt{2}, \sqrt{2}, 0$

Ans: (b)

Solution: $|A - \lambda I| = 0 \Rightarrow \begin{vmatrix} -\lambda & 1 & 0 \\ 1 & -\lambda & 1 \\ 0 & 1 & -\lambda \end{vmatrix} = 0 \Rightarrow -\lambda(\lambda^2 - 1) + \lambda = 0 \Rightarrow \lambda = 0, +\sqrt{2}, -\sqrt{2}$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

GATE-2013

Q17. If \vec{A} and \vec{B} are constant vectors, then $\vec{\nabla}(\vec{A} \cdot (\vec{B} \times \vec{r}))$ is

- (a) $\vec{A} \cdot \vec{B}$ (b) $\vec{A} \times \vec{B}$ (c) \vec{r} (d) zero

Ans: (d)

Solution: Let $\vec{A} = A_0(\hat{x} + \hat{y} + \hat{z})$, $\vec{B} = B_0(\hat{x} + \hat{y} + \hat{z})$ and $\vec{r} = x\hat{x} + y\hat{y} + z\hat{z}$.

$$\vec{B} \times \vec{r} = \hat{x}(z-y)B_0 - \hat{y}(z-x)B_0 + \hat{z}(y-x)B_0 \Rightarrow \vec{\nabla}(\vec{A} \cdot (\vec{B} \times \vec{r})) = 0.$$

Q18. For the function $f(z) = \frac{16z}{(z+3)(z-1)^2}$, the residue at the pole $z=1$ is (your answer should be an integer) _____.

Ans: 3

Solution: At $z=1$ pole is of order 2 so residue is $\frac{1}{2-1} \frac{d^{2-1}}{dz^{2-1}} \left[\frac{(z-1)^2 16z}{(z+3)(z-1)^2} \right]_{z=1} = 3$

Q19. The degenerate eigenvalue of the matrix $\begin{bmatrix} 4 & -1 & -1 \\ -1 & 4 & -1 \\ -1 & -1 & 4 \end{bmatrix}$ is (your answer should be an integer) _____.

Ans: 2, 5, 5

$$\begin{bmatrix} 4-\lambda & -1 & -1 \\ -1 & 4-\lambda & -1 \\ -1 & -1 & 4-\lambda \end{bmatrix} \Rightarrow (4-\lambda) \begin{bmatrix} 1 & -1 & -1 \\ 0 & 5-\lambda & 0 \\ 0 & 0 & 5-\lambda \end{bmatrix} = (4-\lambda)(5-\lambda)^2 = 0 \Rightarrow \lambda = 2, 5, 5.$$

Q20. The number of distinct ways of placing four indistinguishable balls into five distinguishable boxes is _____.

Ans: 120

Solution: $4 \times C_4^5 = 120$ ways

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

GATE-2014

Q21. The unit vector perpendicular to the surface $x^2 + y^2 + z^2 = 3$ at the point $(1, 1, 1)$ is

- (a) $\frac{\hat{x} + \hat{y} - \hat{z}}{\sqrt{3}}$ (b) $\frac{\hat{x} - \hat{y} - \hat{z}}{\sqrt{3}}$ (c) $\frac{\hat{x} - \hat{y} + \hat{z}}{\sqrt{3}}$ (d) $\frac{\hat{x} + \hat{y} + \hat{z}}{\sqrt{3}}$

Ans: (d)

Solution: Let $f = x^2 + y^2 + z^2 - 3 = 0 \Rightarrow \vec{\nabla} f = 2x\hat{x} + 2y\hat{y} + 2z\hat{z}$

$$\Rightarrow \hat{n} = \frac{\vec{\nabla} f}{|\vec{\nabla} f|} \text{ at } (1,1,1) = \frac{2\hat{x} + 2\hat{y} + 2\hat{z}}{\sqrt{12}} = \frac{\hat{x} + \hat{y} + \hat{z}}{\sqrt{3}}$$

Q22. The matrix

$$A = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1+i \\ 1-i & -1 \end{bmatrix} \text{ is}$$

- (a) orthogonal (b) symmetric (c) anti-symmetric (d) Unitary

unitary

unitary $A^\dagger A = I$

Q23. The value of the integral

$$\oint_C \frac{z^2}{e^z + 1} dz$$

where C is the circle $|z| = 4$, is

- (a) $2\pi i$ (b) $2\pi^2 i$ (c) $4\pi^3 i$ (d) $4\pi^2 i$

Ans. (c)

Solution: Pole $e^z = -1$

$$\Rightarrow e^z = e^{i(2m+1)\pi} \text{ where } m = 0, 1, 2, 3, \dots$$

$$\text{Residue Res}_{z=i\pi} = \frac{\phi(z)}{\phi'(z)} = -\frac{\pi^2}{e^{i\pi}} = \pi^2$$

Similarly, $z = -i\pi$, $\text{Res} = \pi^2$

$$\therefore I = 2\pi i (\pi^2 + \pi^2) = 4\pi^3 i$$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

Q24. The solution of the differential equation $\frac{d^2 y}{dt^2} - y = 0$, subject to the boundary conditions

$$y(0) = 1 \text{ and } y(\infty) = 0 \text{ is}$$

(a) $\cos t + \sin t$

(b) $\cosh t + \sinh t$

(c) $\cos t - \sin t$

(d) $\cosh t - \sinh t$

Ans: (d)

Solution:

$$D^2 - 1 = 0 \Rightarrow D = \pm 1 \Rightarrow y(t) = c_1 e^t + c_2 e^{-t}$$

Applying boundary condition

$$y(0) = 1 \Rightarrow 1 = c_1 + c_2 \text{ and } y(\infty) = 0 \Rightarrow 0 = c_1 e^\infty + c_2 e^{-\infty}$$

$$\Rightarrow c_1 = 0, c_2 = 1$$

$$\Rightarrow y(t) = e^{-t} \Rightarrow y(t) = \cosh t - \sinh t$$

GATE-2015

Q25. Consider a complex function $f(z) = \frac{1}{z \left(z + \frac{1}{z} \right) \cos(zx)}$. Which one of the following statements is correct?

(a) $f(z)$ has simple poles at $z = 0$ and $z = -\frac{1}{2}$

(b) $f(z)$ has second order pole at $z = -\frac{1}{2}$

(c) $f(z)$ has infinite number of second order poles

(d) $f(z)$ has all simple poles

Ans.: (b)

Solution:
$$f(z) = \frac{1}{z \left(z + \frac{1}{z} \right) \cos(z\pi)}$$

For k^{th} order pole $\because \lim_{z \rightarrow a} (z - a)^n f(z) = \text{finite and } \neq 0$

At $z = 0$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

$\lim_{z \rightarrow 0} zf'(z) = \text{finite}$ $z = 0$ is a simple pole.

At $z = -\frac{1}{2}$

$$\begin{aligned} \lim_{z \rightarrow -\frac{1}{2}} \frac{\left(z + \frac{1}{2}\right)^2}{z \left(z + \frac{1}{2}\right) \cos z\pi} &= \lim_{z \rightarrow -\frac{1}{2}} \frac{\left(z + \frac{1}{2}\right)}{z \cos z\pi} = \lim_{z \rightarrow -\frac{1}{2}} \frac{1}{1 \cdot \cos z\pi + z \cdot \pi (-\sin z\pi)} \\ &= \lim_{z \rightarrow -\frac{1}{2}} \frac{1}{\cos z\pi - z\pi \sin z\pi} = \frac{1}{\frac{\pi}{2}} = \frac{2}{\pi} = \text{finite} \end{aligned}$$

$\Rightarrow f(z)$ has second order pole at $z = -\frac{1}{2}$

Q26. The value of $\int_0^3 t^2 \delta(3t-6) dt$ is _____ (upto one decimal place)

Ans.: 1.33

Solution: $\int_0^3 t^2 \delta(3t-6) dt = \int_0^3 t^2 \delta[3(t-2)] dt = \frac{1}{3} \int_0^3 t^2 \delta(t-2) dt = \frac{4}{3}$

Q27. If $f(x) = e^{-x^2}$ and $g(x) = |x|e^{-x^2}$, then

- (a) f and g are differentiable everywhere
- (b) f is differentiable everywhere but g is not
- (c) g is differentiable everywhere but f is not
- (d) g is discontinuous at $x = 0$

Ans. (b)

Solution: $f(x) = e^{-x^2}$ is differentiable but $g(x) = |x|e^{-x^2}$ is not differentiable.

$$g(x) = \begin{cases} -xe^{-x^2}; & x < 0 \\ xe^{-x^2}; & x > 0 \end{cases}$$

Left hand Limit $\lim_{h \rightarrow 0} g(x-h) = -(x-h)e^{-(x-h)^2}$

Right hand Limit $\lim_{h \rightarrow 0} g(x+h) = (x+h)e^{-(x+h)^2}$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

$$\Rightarrow \lim_{h \rightarrow 0} g(x-h) \neq \lim_{h \rightarrow 0} g(x+h)$$

Q28. Consider $w = f(z) = u(x, y) + iv(x, y)$ to be an analytic function in a domain D . Which one of the following options is NOT correct?

(a) $u(x, y)$ satisfies Laplace equation in D

(b) $v(x, y)$ satisfies Laplace equation in D

(c) $\int_{z_1}^{z_2} f(z) dz$ is dependent on the choice of the contour between z_1 and z_2 in D

(d) $f(z)$ can be Taylor expanded in D

Ans.: (c)

Solution: $w = f(z) = u(x, y) + iv(x, y)$ to be an analytic function in a domain D , $\int_{z_1}^{z_2} f(z) dz$ is independent on the choice of the contour between z_1 and z_2 in D .

Q29. The Heaviside function is defined as $H(t) = \begin{cases} +1 & \text{for } t > 0 \\ -1 & \text{for } t < 0 \end{cases}$ and its Fourier transform is

given by $-2i/\omega$. The Fourier transform of $\frac{1}{2}[H(t+1/2) - H(t-1/2)]$ is

(a) $\frac{\sin\left(\frac{\omega}{2}\right)}{\frac{\omega}{2}}$

(b) $\frac{\cos\left(\frac{\omega}{2}\right)}{\frac{\omega}{2}}$

(c) $\sin\left(\frac{\omega}{2}\right)$

(d) 0

Ans.: (a)

Solution: $H(f) = \int_{-\infty}^{\infty} h(t) e^{-i2\pi ft} dt$

For a function $h(t)$, $H(f) = -\frac{2i}{\omega}$

For $h(t-t_0)$, Fourier Transform is $e^{-i2\pi ft_0} H(f)$

Shifting theorem

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

$$\text{For } \frac{1}{2} \left[h\left(t + \frac{1}{2}\right) - h\left(t - \frac{1}{2}\right) \right] = \frac{1}{2} \left[e^{i\frac{\omega}{2}} - e^{-i\frac{\omega}{2}} \right] \frac{-2i}{\omega} = \frac{1}{2i} \left[e^{i\frac{\omega}{2}} - e^{-i\frac{\omega}{2}} \right] \frac{-2i}{\omega} \times i$$

$$\text{The Fourier transform of } \frac{1}{2} \left[H\left(t + \frac{1}{2}\right) - H\left(t - \frac{1}{2}\right) \right] = \frac{\sin\left(\frac{\omega}{2}\right)}{\frac{\omega}{2}} .$$

Q30. A function $y(z)$ satisfies the ordinary differential equation $y'' + \frac{1}{z}y' - \frac{m^2}{z^2}y = 0$, where $m = 0, 1, 2, 3, \dots$. Consider the four statements P, Q, R, S as given below.

P: z^m and z^{-m} are linearly independent solutions for all values of m

Q: z^m and z^{-m} are linearly independent solutions for all values of $m > 0$

R: $\ln z$ and 1 are linearly independent solutions for $m = 0$

S: z^m and $\ln z$ are linearly independent solutions for all values of m

The correct option for the combination of valid statements is

- (a) P, R and S only (b) P and R only (c) Q and R only (d) R and S only

Ans.: (c)

$$\text{Solution: } y'' + \frac{1}{z}y' - \frac{m^2}{z^2}y = 0 \Rightarrow z^2y'' + zy' - m^2y = 0$$

$$m = 0, 1, 2, 3, \dots \quad z = e^x, \quad D = \frac{d}{dx}$$

$$\text{If } m = 0; \quad z^2y'' + zy' = 0$$

$$\left[D(D-1) + D \right] y = 0 \Rightarrow \left[D^2 - D + D \right] y = 0 \Rightarrow (D^2 - m^2)y = 0$$

$$y = c_1 + c_2x$$

$$D = \pm m$$

$$y = c_1 + c_2 \ln z$$

$$c_1 e^{mx} + c_2 e^{-mx}$$

R is correct.

$$c_1 e^{m \log z} + c_2 e^{-m \log z}$$

or if $m \neq 0, m > 0$

$$y = c_1 \cosh(m \log(z)) + ic_2 \sinh(m \log(x)) \Rightarrow m > 0$$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

GATE-2016

Q31. Consider the linear differential equation $\frac{dy}{dx} = xy$. If $y = 2$ at $x = 0$, then the value of y at

$x = 2$ is given by

- (a) e^{-2} (b) $2e^{-2}$ (c) e^2 (d) $2e^2$

Ans.: (d)

Solution: $\frac{dy}{dx} = xy \Rightarrow \frac{1}{y} dy = x dx \Rightarrow \ln y = \frac{x^2}{2} + \ln c \Rightarrow y = ce^{x^2/2}$

If $y = 2$ at $x = 0 \Rightarrow c = 2 \Rightarrow y = 2e^{x^2/2}$.

The value of y at $x = 2$ is given by $y = 2e^2$

Q32. Which of the following is an analytic function of z everywhere in the complex plane?

- (a) z^2 (b) $(z^*)^2$ (c) $|z|^2$ (d) \sqrt{z}

Ans.: (a)

Solution: $z^2 = (x + iy)^2 = x^2 - y^2 + i(2xy) \Rightarrow u = x^2 - y^2$ and $v = 2xy$

Cauchy Riemann equations $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = 2x$, $\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} = 2y$ satisfies.

Q33. The direction of $\vec{\nabla}f$ for a scalar field $f(x, y, z) = \frac{1}{2}x^2 - xy + \frac{1}{2}z^2$ at the point $P(1,1,2)$ is

- (a) $\frac{(-\hat{j} - 2\hat{k})}{\sqrt{5}}$ (b) $\frac{(-\hat{j} + 2\hat{k})}{\sqrt{5}}$ (c) $\frac{(\hat{j} - 2\hat{k})}{\sqrt{5}}$ (d) $\frac{(\hat{j} + 2\hat{k})}{\sqrt{5}}$

Ans.: (b)

Solution: $\vec{\nabla}f = (x - y)\hat{i} - \hat{x}\hat{j} + z\hat{k} \Rightarrow \hat{n} = \left(\frac{\vec{\nabla}f}{|\vec{\nabla}f|} \right)_{1,1,2} = \frac{-\hat{j} + 2\hat{k}}{\sqrt{5}}$

Q34. A periodic function $f(x)$ of period 2π is defined in the interval $(-\pi < x < \pi)$

$$f(x) = \begin{cases} -1, & -\pi < x < 0 \\ 1, & 0 < x < \pi \end{cases}$$

The appropriate Fourier series expansion for $f(x)$ is

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

$$(a) f(x) = \left(\frac{4}{\pi}\right) [\sin x + (\sin 3x)/3 + (\sin 5x)/5 + \dots]$$

$$(b) f(x) = \left(\frac{4}{\pi}\right) [\sin x - (\sin 3x)/3 + (\sin 5x)/5 - \dots]$$

$$(c) f(x) = \left(\frac{4}{\pi}\right) [\cos x + (\cos 3x)/3 + (\cos 5x)/5 + \dots]$$

$$(d) f(x) = \left(\frac{4}{\pi}\right) [\cos x - (\cos 3x)/3 + (\cos 5x)/5 - \dots]$$

Ans.: (a)

$$\text{Solution: } f(x) = \begin{cases} -1, & -\pi < x < 0 \\ 1, & 0 < x < \pi \end{cases}$$

$$\text{Let } f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$\therefore a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$\Rightarrow a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \left[\int_{-\pi}^0 (-1) dx + \int_0^{\pi} (1) dx \right] = \frac{1}{2\pi} \left[[-x]_{-\pi}^0 + [x]_0^{\pi} \right] = 0$$

This can also be seen without integration, since the area under the curve of $f(x)$ between $-\pi$ to π is zero.

$$\therefore a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nxdx$$

$$\Rightarrow a_n = \frac{1}{\pi} \left[\int_{-\pi}^0 (-1) \cos nxdx + \int_0^{\pi} (1) \cos nxdx \right] = \frac{1}{\pi} \left[-\left\{ \frac{\sin nx}{n} \right\}_{-\pi}^0 + \left\{ \frac{\sin nx}{n} \right\}_0^{\pi} \right] = 0$$

$$\therefore b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nxdx$$

$$\Rightarrow b_n = \frac{1}{\pi} \left[\int_{-\pi}^0 (-1) \sin nxdx + \int_0^{\pi} (1) \sin nxdx \right]$$

$$\Rightarrow b_n = \frac{1}{\pi} \left[\left\{ \frac{\cos nx}{n} \right\}_{-\pi}^0 - \left\{ \frac{\cos nx}{n} \right\}_0^{\pi} \right] = \frac{1}{\pi} \left[\frac{1}{n} - \frac{(-1)^n}{n} - \frac{(-1)^n}{n} + \frac{1}{n} \right] = \frac{1}{\pi} \left[\frac{2}{n} - \frac{2(-1)^n}{n} \right]$$

Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16

If n is even $b_n = 0$ and If n is odd $b_n = \frac{4}{n\pi}$.

Thus Fourier series is $f(x) = \frac{4}{\pi} \left[\sin x + \frac{1}{3} \sin 3x + \frac{1}{5} \sin 5x + \dots \right]$



Head office

fiziks, H.No. 23, G.F, Jia Sarai,
Near IIT, Hauz Khas, New Delhi-16
Phone: 011-26865455/+91-9871145498

Branch office

Anand Institute of Mathematics,
28-B/6, Jia Sarai, Near IIT
Hauz Khas, New Delhi-16